

STM - Structure Search  
5-19-04

2 inventors d ibib abs hitstr 1-10



L5 ANSWER 1 OF 10 CAPLUS COPYRIGHT 2004 ACS on STN  
 ACCESSION NUMBER: 2004:80450 CAPLUS  
 DOCUMENT NUMBER: 140:145835  
 TITLE: Preparation of dibenzofused bicyclo[2.2.2]octane-derived amides as modulators of the glucocorticoid receptor  
 INVENTOR(S): Vaccaro, Wayne; Yang, Bingwei Vera; Kim, Soong-hoon; Huynh, Tram; Tortolani, David R.; Leavitt, Kenneth J.; Li, Wenying; Doweiko, Arthur M.; Chen, Xiao-tao; Doweiko, Lidia  
 PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA; et al.  
 SOURCE: PCT Int. Appl., 265 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004009017	A2	20040129	WO 2003-US22300	20030717
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

PRIORITY APPLN. INFO.: US 2002-396877P P 20020718


OTHER SOURCE(S): MARPAT 140:145835

GI



AB Title compds. I [R-R4 = H, alk(en/yn)yl, alkoxy, aryl, etc.; Z = carboxamido, alkylamino, etc.] are prepared For instance, 2-amino-4,5-dimethylthiazole is coupled to the acid derived from the cycloaddn. of methacrylic acid and anthracene (CH3CN, EDCI, Et3N, HOAt, 18 h) to give II. I are glucocorticoid receptor modulators which are useful in treating diseases requiring glucocorticoid receptor agonist or antagonist therapy such as obesity, diabetes, inflammatory and immune disorders.

IT 312317-98-9P 650625-43-7P 650625-55-1P



L5 ANSWER 2 OF 10 CAPLUS COPYRIGHT 2004 ACS on STN  
 ACCESSION NUMBER: 2004:80449 CAPLUS  
 DOCUMENT NUMBER: 140:157927  
 TITLE: Homology modeling of nuclear hormone receptor Site II  
 and design of Site II ligands  
 INVENTOR(S): Dowejko, Arthur; Nadler, Steven G.  
 PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA  
 SOURCE: PCT Int. Appl., 276 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004009016	A2	20040129	WO 2003-US22299	20030717

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,  
 CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
 GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
 LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM,  
 PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN,  
 TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY,  
 KG, KZ, MD, RU

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,  
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,  
 NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
 GW, ML, MR, NE, SN, TD, TG


PRIORITY APPLN. INFO.: US 2002-396907P R 20020718

AB A binding site in nuclear hormone receptors is described and its  
 structural coordinates are provided. The invention provides  
 machine-readable data storage media comprising structure coordinates of  
 Site II and computer systems comprising the machine-readable data storage  
 media. The invention provides methods used in the design and  
 identification of ligands of Site II and of modulators of nuclear hormone  
 receptors. The invention provides ligands of Site II, modulators of NHRs,  
 pharmaceutical comps. comprising modulators of NHRs, methods of  
 modulating NHRs, and methods of treating diseases by administering  
 modulators of an NHR. Also provided are methods of designing mutants,  
 mutant NHRs, Site II binding assays, and models of Site II.

IT 312317-98-9P 650625-43-7P 650625-47-1P  
 650625-55-1P 650625-58-4P 650625-60-8P  
 650625-62-0P 650625-64-2P 650625-67-5P  
 650625-68-6P 650625-70-0P 650625-72-2P  
 650625-74-4P 650625-75-5P 650625-76-6P

*Same  
 priority  
 date*

Absolute stereochemistry.



L5 ANSWER 3 OF 10 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:501478 CAPLUS

DOCUMENT NUMBER: 137:229148

TITLE: In vitro increase in chloroquine accumulation induced by dihydroethano- and ethenoanthracene derivatives in Plasmodium falciparum-parasitized erythrocytes

AUTHOR(S): Pradines, Bruno; Alibert, Sandrine; Houdoin, Carole; Santelli-Rouvier, Christiane; Mosnier, Joel; Fusai, Thierry; Rogier, Christophe; Barbe, Jacques; Parzy, Daniel

CORPORATE SOURCE: Unite de Parasitologie, Institut de Medecine Tropicale du Service de Sante des Armees, Institut Federatif de la Recherche 48, Marseille, 13998, Fr.

SOURCE: Antimicrobial Agents and Chemotherapy (2002), 46(7), 2061-2068

CODEN: AMACCQ; ISSN: 0066-4804

PUBLISHER: American Society for Microbiology

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effects of a series of dihydroethano- and ethenoanthracene derivs. on chloroquine (CQ) accumulation in CQ-susceptible strain 3D7 and CQ-resistant clone W2 were assessed. The levels of CQ accumulation increased little or none in CQ-susceptible strain 3D7 and generally increased markedly in CQ-resistant strain W2. At 10  $\mu$ M, 28 compds. yielded cellular accumulation ratios (CARs) greater than that observed with CQ alone in W2. At 10  $\mu$ M, in strain W2, 21 of 31 compds. had CQ CARs two or more times higher than that of CQ alone, 15 of 31 compds. had CQ CARs three or more times higher than that of CQ alone, 13 of 31 compds. had CQ CARs four or more times higher than that of CQ alone, and 9 of 31 compds. had CQ CARs five or more times higher than that of CQ alone. At 1  $\mu$ M, 17 of 31 compds. had CQ CARs two or more times higher than that of CQ alone, 12 of 31 compds. had CQ CARs three or more times higher than that of CQ alone, 6 of 31 compds. had CQ CARs four or more times higher than that of CQ alone, and 3 of 31 compds. had CQ CARs five or more times higher than that of CQ alone. At 1  $\mu$ M, 17 of 31 compds. were more potent inducers of CQ accumulation than verapamil and 12 of 31 compds. were more potent inducers of CQ accumulation than promethazine. The nature of the basic group seems to be associated with increases in the levels of CQ accumulation. At 1 and 10  $\mu$ M, 10 of 14 and 13 of 14 compds. with amino group (amines and diamines), resp., had CARs  $\geq 3$ , while at 1 and 10  $\mu$ M, only 1 of the 13 derivs. with amido groups had CARs  $\geq 3$ . Among 12 of the 31 compds. which were more active inducers of CQ accumulation than promethazine at 1  $\mu$ M, 10 had amino groups and 1 had an amido group.

IT 448958-12-1, BG 1050 448958-13-2, BG 1051

448958-14-3, BG 1049


10/621,909

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(in vitro increase in chloroquine accumulation induced by  
dihydroethano- and ethenoanthracene derivs. in Plasmodium  
falciparum-parasitized erythrocytes)

RN 448958-12-1 CAPLUS

CN 9,10-Ethanoanthracene-11-carboxamide, 9,10-dihydro-12-methyl-N-2-pyridinyl-  
, (11R,12S)-rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 448958-13-2 CAPLUS

CN 9,10-Ethanoanthracene-11-carboxamide, 9,10-dihydro-12-methyl-N-3-pyridinyl-  
, (11R,12S)-rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 448958-14-3 CAPLUS

CN 9,10-Ethanoanthracene-11-carboxamide, 9,10-dihydro-12-methyl-N-4-pyridinyl-  
, (11R,12S)-rel- (9CI) (CA INDEX NAME)

Relative stereochemistry.



REFERENCE COUNT: 60 THERE ARE 60 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 4 OF 10 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 2002:462893 CAPLUS

10/621,909

DOCUMENT NUMBER: 137:185297  
TITLE: Synthesis and Effects on Chloroquine Susceptibility in Plasmodium falciparum of a Series of New Dihydroanthracene Derivatives  
AUTHOR(S): Alibert, Sandrine; Santelli-Rouvier, Christiane; Pradines, Bruno; Houdoin, Carole; Parzy, Daniel; Karolak-Wojciechowska, Janina; Barbe, Jacques  
CORPORATE SOURCE: GERCTOP-UMR CNRS 6009, Faculte de Pharmacie, Universite de la Mediterranee, Marseille, 13385, Fr.  
SOURCE: Journal of Medicinal Chemistry (2002), 45(15), 3195-3209  
CODEN: JMCMAR; ISSN: 0022-2623  
PUBLISHER: American Chemical Society  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
OTHER SOURCE(S): CASREACT 137:185297


AB To suggest a mechanism of action for drugs capable of reversing the chloroquine resistance in Plasmodium falciparum, a new set of 9,10-dihydro-9,10-ethano- and -ethenoanthracene derivs. was synthesized and tested to assess their effect on chloroquine susceptibility in resistant strains of Plasmodium falciparum. With respect to this, reversal of resistance and change in drug accumulation were compared. Structure-activity relationship and mol. modeling studies made it possible to define a pharmacophore moiety for reversal agents and to propose a putative model of interaction with some selected amino acids.

IT 448958-12-1P 448958-13-2P 448958-14-3P  
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(preparation and effects on chloroquine susceptibility in Plasmodium falciparum of a series of ethano- and ethenodihydroanthracenes)

RN 448958-12-1 CAPLUS

CN 9,10-Ethanoanthracene-11-carboxamide, 9,10-dihydro-12-methyl-N-2-pyridinyl-, (11R,12S)-rel- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 448958-13-2 CAPLUS

CN 9,10-Ethanoanthracene-11-carboxamide, 9,10-dihydro-12-methyl-N-3-pyridinyl-, (11R,12S)-rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.

10/621,909



RN 448958-14-3 CAPLUS  
CN 9,10-Ethanoanthracene-11-carboxamide, 9,10-dihydro-12-methyl-N-4-pyridinyl-, (11R,12S)-rel- (9CI) (CA INDEX NAME)

Relative stereochemistry.



REFERENCE COUNT: 48 THERE ARE 48 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 5 OF 10 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:220218 CAPLUS

DOCUMENT NUMBER: 130:237374

TITLE: New tetracyclo[6.6.2.02,7.09,14]hexadeca-2(7),3,5,9(14),10,12]hexaenes as phospholipase inhibitors

INVENTOR(S): Friebe, Walter-Gunar; Tibes, Ulrich; Scheuer, Werner

PATENT ASSIGNEE(S): Roche Diagnostics G.m.b.H., Germany

SOURCE: Ger. Offen., 10 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19742014	A1	19990325	DE 1997-19742014	19970924
ZA 9808711	A	20000323	ZA 1998-8711	19980923
CA 2304879	AA	19990401	CA 1998-2304879	19980924
WO 9915493	A1	19990401	WO 1998-EP6096	19980924

W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

10/621,909

AU 9897466	A1	19990412	AU 1998-97466	19980924
BR 9813217	A	20000829	BR 1998-13217	19980924
EP 1034162	A1	20000913	EP 1998-951464	19980924
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI				
TR 200001222	T2	20000921	TR 2000-200001222	19980924
JP 2002505999	T2	20020226	JP 2000-512804	19980924
PRIORITY APPLN. INFO.:			DE 1997-19742014 A	19970924
			WO 1998-EP6096	W 19980924
OTHER SOURCE(S):		MARPAT 130:237374		
GI				




AB Title compds. I [R1, R2 = H, halogen; X = H, Y = (un)substituted NH2, N+H2Me; XY = (un)substituted CH2NH; Z = CH2, C:NH] were prepared for use as phospholipase inhibitors (no data). Thus, I [R1, R2, X = H, Y = NH2.HCl, Z = CH2] was treated with KOCN to give I [R1, R2, X = H, Y = NHCONH2, Z = CH2].

IT **221352-72-3P**  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation of tetracyclohexadecahexaenes as phospholipase inhibitors)

RN 221352-72-3 CAPLUS

CN 9,10-Ethanoanthracene-11-carboxamide, 9,10-dihydro-N-4-pyridinyl- (9CI)  
(CA INDEX NAME)




IT **221352-19-8P 221352-21-2P 221352-47-2P**  
**221352-50-7P 221352-54-1P**  
RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation of tetracyclohexadecahexaenes as phospholipase inhibitors)

RN 221352-19-8 CAPLUS


CN 4-Pyridinamine, N-[(9,10-dihydro-9,10-ethanoanthracen-11-yl)methyl]- (9CI)  
(CA INDEX NAME)

10/621,909




RN 221352-21-2 CAPLUS

CN 4-Pyridinamine, N-[(1,5-dichloro-9,10-dihydro-9,10-ethanoanthracen-11-yl)methyl]- (9CI) (CA INDEX NAME)




RN 221352-47-2 CAPLUS

CN 1H-Imidazol-2-amine, N-[(1,5-dichloro-9,10-dihydro-9,10-ethanoanthracen-11-yl)methyl]-4,5-dihydro- (9CI) (CA INDEX NAME)



RN 221352-50-7 CAPLUS


CN 1H-Imidazol-2-amine, N-[(9,10-dihydro-9,10-ethanoanthracen-11-yl)methyl]-4,5-dihydro- (9CI) (CA INDEX NAME)



RN 221352-54-1 CAPLUS

CN 1H-Imidazol-2-amine, N-[(4,5-dichloro-9,10-dihydro-9,10-ethanoanthracen-11-yl)methyl]-4,5-dihydro-, monohydrobromide (9CI) (CA INDEX NAME)





● HBr

L5 ANSWER 6 OF 10 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1995:457943 CAPLUS

DOCUMENT NUMBER: 123:199069

TITLE: Infrared spectra of N-ferrocenylamic acids

AUTHOR(S): Yang, Bingqin; Ma Huairang; Chen Zhibin

CORPORATE SOURCE: Department of Chemistry, Northwest University, Xian, 710069, Peop. Rep. China

SOURCE: Guangpuxue Yu Guangpu Fenxi (1995), 15(1), 53-6  
CODEN: GYGFE; ISSN: 1000-0593

PUBLISHER: Beijing Daxue Chubanshe

DOCUMENT TYPE: Journal

LANGUAGE: Chinese

AB The IR spectra of seven N-ferrocenylamic acids are reported. The effect of intramol. hydrogen bonds was discussed and the absorption characteristics of some structural isomers were reported. In addition, the mol. structures of some interconversion isomers were proven by IR.

IT 146083-59-2


RL: PRP (Properties)


(IR spectra of N-ferrocenylamic acids)

RN 146083-59-2 CAPLUS

CN Ferrocene, [[(12-carboxy-9,10-dihydro-9,10-ethanoanthracen-11-yl)carbonyl]amino]-, cis- (9CI) (CA INDEX NAME)

PAGE 1-A





L5 ANSWER 7 OF 10 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1994:244394 CAPLUS

DOCUMENT NUMBER: 120:244394

TITLE: Dibenzo-fused derivatives of bicyclo[2.2.2]octane as cholecystokinin inhibitors

INVENTOR(S): Kalindjian, Sarkis Barret; Low, Caroline Minli Rachel; McDonald, Iain Mair; Hull, Robert Antony David; Shankley, Nigel Paul; Buck, Ildiko Maria; Steel, Katherine Isobel Mary; Davies, Jonathan Michael Richar; Dunstone, David John; et al.

PATENT ASSIGNEE(S): James Black Foundation Ltd., UK

SOURCE: PCT Int. Appl., 159 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English



FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9316982	A1	19930902	WO 1993-GB346	19930219
W: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG				
AU 9335097	A1	19930913	AU 1993-35097	19930219
ZA 9301193	A	19940819	ZA 1993-1193	19930219
EP 626942	A1	19941207	EP 1993-904230	19930219
EP 626942	B1	19970423		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
JP 07504184	T2	19950511	JP 1993-514633	19930219
HU 71499	A2	19951128	HU 1994-2280	19930219
AT 152095	E	19970515	AT 1993-904230	19930219

10/621,909

US 5514683	A	19960507	US 1994-288185	19940809
NO 9403055	A	19941011	NO 1994-3055	19940818
FI 9403817	A	19940819	FI 1994-3817	19940819
PRIORITY APPLN. INFO.:			GB 1992-3608	A 19920220
			GB 1992-13093	A 19920619
			GB 1992-24629	A 19921124
			WO 1993-GB346	A 19930219
			GB 1993-16722	A 19930812
OTHER SOURCE(S):			MARPAT 120:244394	
GI				




- AB Title compds. I [W = CO, SO, SO<sub>2</sub>; X = CO, SO, SO<sub>2</sub>, COCH<sub>2</sub> (with CO end bound to Y), provided that  $\geq 1$  of W and X contains CO; Y = certain (un)substituted OH or NH<sub>2</sub> groups; Z = different (un)substituted OH or NH<sub>2</sub> groups; R<sub>1</sub> = H, Me, halo, (amidated or esterified) CO<sub>2</sub>H or CH<sub>2</sub>CO<sub>2</sub>H; R<sub>2</sub> = groups for R<sub>1</sub>, or COZ' (Z' = Z) when Z is absent and W = H; or R<sub>1</sub>R<sub>2</sub> = pi bond; R<sub>3</sub>, R<sub>4</sub> = halo, amino, NO<sub>2</sub>, cyano, SO<sub>2</sub>NH<sub>2</sub>, alkyl, alkoxy, (amidated or esterified) CO<sub>2</sub>H; R<sub>5</sub>, R<sub>6</sub> = H, R<sub>3</sub>; m, n = 0-4, provided that both are  $\leq 2$  unless R<sub>3</sub> or R<sub>4</sub>, resp., are exclusively halo] were prepared as ligands binding at cholecystokinin (CCK) and gastrin receptors. Thus, 2,3,5,6-dibenzobicyclo[2.2.2]octane-7,8-dicarboxylic acid anhydride reacted with 1-adamantanemethylamine, the resultant acid-amide was condensed with H-L-Pro-OCH<sub>2</sub>Ph.HCl using PyBOP, and the benzyl ester function was hydrogenolyzed and reesterified with diazomethane to give title compound cis-II as a mixture of 2 diastereomers which were separated by repeated crystallization. These isomers bound to CCKB receptors (mouse cortical membrane) with pK<sub>i</sub> = 5.8 and 7.3. Included are 238 synthetic examples, 1H NMR data for all final products (free bases or N-methyl-D-glucamine salts), and receptor-binding results (CCKA, CCKB, and gastrin) for most I.
- IT **153459-03-1P**  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(preparation of, as CCK and gastrin antagonist)
- RN 153459-03-1 CAPLUS
- CN 2-Furancarboxylic acid, 5-[[[9,10-dihydro-12-[[[tricyclo[3.3.1.1<sup>3,7</sup>]dec-1-ylmethyl]amino]carbonyl]-9,10-ethanoanthracen-11-yl]carbonyl]amino]-,


10/621,909

methyl ester, cis- (9CI) (CA INDEX NAME)

Relative stereochemistry.



L5 ANSWER 8 OF 10 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 1993:102173 CAPLUS  
DOCUMENT NUMBER: 118:102173  
TITLE: Synthesis and reaction of N-ferrocenylamic acid  
AUTHOR(S): Ma, Huairang; Chen, Zhibing; Yang, Bingqin; Li, Jinsong  
CORPORATE SOURCE: Dep. Chem., Northwest Univ., Xian, 710069, Peop. Rep. China  
SOURCE: Gaodeng Xuexiao Huaxue Xuebao (1992), 13(5), 633-5  
CODEN: KTHPDM; ISSN: 0251-0790  
DOCUMENT TYPE: Journal  
LANGUAGE: Chinese  
GI




AB Seven N-ferrocenylamic acids have been synthesized and characterized by elemental anal., IR and <sup>1</sup>H NMR. The investigation on the chelation of acid I (Fc = ferrocenyl) with Cu<sup>2+</sup> and Zn<sup>2+</sup> ions demonstrates the coordination of the oxygen atom on the amide group to the metallic ion.

IT **146083-59-2P**  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)


RN 146083-59-2 CAPLUS

CN Ferrocene, [[[12-carboxy-9,10-dihydro-9,10-ethanoanthracen-11-yl)carbonyl]amino]-, cis- (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A



L5 ANSWER 9 OF 10 CAPLUS COPYRIGHT 2004 ACS on STN  
 ACCESSION NUMBER: 1989:149699 CAPLUS  
 DOCUMENT NUMBER: 110:149699  
 TITLE: Solid precursors, methods, and apparatus for  
 development of latent fingerprints with cyanoacrylates  
 INVENTOR(S): Warrener, Ronald Norman; Yong, Siaw Jan  
 PATENT ASSIGNEE(S): Australian National University, Australia  
 SOURCE: PCT Int. Appl., 50 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 8801616	A1	19880310	WO 1987-AU286	19870825
W: AU, JP, US				
RW: AT, BE, CH, DE, FR, GB, IT, LU, NL, SE				

AU 8779188            A1    19880324            AU 1987-79188            19870825  
 PRIORITY APPLN. INFO.:            AU 1986-7669            19860825  
    WO 1987-AU286            19870825


AB    Solid precursors of cyanoacrylate monomers, such as Diels-Alder adducts, homopolymers, or cyclic or acyclic precursors, may be used in a method for the development of a latent fingerprint on a surface by exposure of the surface to cyanoacrylate monomer generated from a solid precursor to form a cyanoacrylate-developed print. Preferably, the solid precursor includes colored or fluorescent substituent groups. An apparatus for use in the cyanoacrylate development of latent fingerprints is disclosed also. A mixture of anthracene, Super Glue, hydroquinone, and dry benzene was refluxed at 90° for 12 h. The anthracene/Et 2-cyanoacrylate adduct was treated with NaOH to make anthracene/2-cyanoacrylic acid adduct (m. 206-209°) in 99.7% yield.

IT    **119858-97-8**

RL: PRP (Properties)  
 (NMR of)

RN    119858-97-8    CAPLUS

CN    9,10-Ethanoanthracene-11-carboxamide, 11-cyano-9,10-dihydro-N-8-quinolinyl-  
 (9CI)    (CA INDEX NAME)




IT    **119859-04-0 119859-05-1**

RL: BIOL (Biological study)  
 (as solid precursor for cyanoacrylate development of latent fingerprints)

RN    119859-04-0    CAPLUS


CN    9,10-Ethanoanthracene-11-carboxamide, 11-cyano-9,10-dihydro-N-(4-methyl-2-oxo-2H-1-benzopyran-7-yl)- (9CI)    (CA INDEX NAME)



RN    119859-05-1    CAPLUS

10/621,909

CN 9,10-Ethanoanthracene-11-carboxamide, 11-cyano-9,10-dihydro-N-(2-oxo-2H-1-benzopyran-7-yl)- (9CI) (CA INDEX NAME)



L5 ANSWER 10 OF 10 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 1979:120619 CAPLUS  
DOCUMENT NUMBER: 90:120619  
TITLE: Reactions of aminopyridines with some inner anhydrides  
AUTHOR(S): El-Zanfally, S.; El-Basil, S.  
CORPORATE SOURCE: Org. Chem. Dep., Fac. Pharm., Cairo, Egypt  
SOURCE: Egyptian Journal of Pharmaceutical Sciences (1978),  
Volume Date 1976, 17(1), 53-62  
CODEN: EJPSBZ; ISSN: 0301-5068

DOCUMENT TYPE: Journal  
LANGUAGE: English


AB Reaction of 3-aminopyridine inner anhydrides, e.g., succinic or maleic anhydride, gives amic acids. In the case of 2- and 4-aminopyridines, the products obtained were amic acids or cyclized products (succinimide derivs.). With maleic anhydride, both 2- and 4-aminopyridines form a charge-transfer complex. Phenylsuccinic anhydride behaves as a C acid with 4- but not with 2-aminopyridine.

IT 69537-47-9P 69537-51-5P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 69537-47-9 CAPLUS


CN 9,10-Ethanoanthracene-11-carboxylic acid, 9,10-dihydro-12-[(2-pyridinylamino)carbonyl]-, cis- (9CI) (CA INDEX NAME)



RN 69537-51-5 CAPLUS

CN 9,10-Ethanoanthracene-11-carboxylic acid, 9,10-dihydro-12-[(4-pyridinylamino)carbonyl]-, cis- (9CI) (CA INDEX NAME)

10/621,909



=> d his

(FILE 'HOME' ENTERED AT 14:10:23 ON 19 MAY 2004)

FILE 'REGISTRY' ENTERED AT 14:10:31 ON 19 MAY 2004

L1 STRUCTURE UPLOADED

L2 25 S L1

L3 588 S L1 FULL

FILE 'CAPLUS' ENTERED AT 14:11:24 ON 19 MAY 2004


L4 2 S L3/THU

L5 10 S L3

=> d l1

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=>